



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

January 31, 2007

Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569

Subject: Label Amendment NAFTA label
EPA Reg. No. 10163-287 Far-Go Granular Herbicide
EPA Receipt date: January 31, 2007

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable.

Amended labeling will supercede all previously accepted ones. A stamped copy of the labeling is enclosed for your records. Submit one (1) copy of final printed labeling before you release the product for shipment. If you have any questions please contact me at 703-305-5697.

All subsequent amendments to this label must be submitted to Canada and the United States at the same time for joint review.

Sincerely,

A handwritten signature in black ink, appearing to read "James Tompkins", is written over a circular stamp. The signature is fluid and cursive.

James Tompkins
Product manager 25
Herbicide Branch
Registration Division (7505P)

IMPORTANT NOTICE

This product is registered in both the U.S. and Canada. It has BOTH U.S. and Canada Labeling. You must follow ONLY the U.S. label WHEN USING THIS product in the U.S. You must follow ONLY the Canadian label WHEN USING THIS product in Canada

Far-GO Granular

In the United States

Herbicide

Avadex®
Micro Active

Herbicide

In Canada

GROUP 8 HERBICIDE

Granular Herbicide for Fall or Spring Treatment in Conventional and Direct Seeding Systems to Control Wild Oats in labeled crops. See the appropriate U.S. or Canadian Use Directions.

AGRICULTURAL

ACTIVE INGREDIENTS/GUARANTEE

Thialate, S-(2,3,3-trichloroallyl)-disopropylthiocarbamate

OTHER INGREDIENTS:

ATTENTION, contains the allergen soy.

% By Wt.
10.0%
90.0%
TOTAL 100.0%

CAUTION

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. Use only according to label instructions. Read "NOTICE OF CONDITIONS OF SALE AND WARRANTY LIMITATIONS" before buying or using. If terms are not acceptable, return at once unopened.

FIRST AID Herbicide

If swallowed	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

When seeking medical attention, take the container label if at all possible. If not, take information which identifies the product, that is, the product names and registration numbers. You may also contact 1-888-478-0798 for emergency medical treatment information.

TOXICOLOGICAL INFORMATION: Treat symptomatically

United States AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40CFR Part 170. Refer to supplemental labeling under "Agricultural Use Requirements." In the Directions for Use section for information about this standard.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

NET CONTENTS 1000 lbs./453.60 kgs

ACCEPTED

EPA Est. No.

Registration No. 25112 PEST CONTROL PRODUCTS ACT

EPA Reg. No. 10163-287

FFG-ADX-NAFTA-01

Goway
The Go To Company

Goway
CANADA

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide Yuma, AZ 85364 U.S.A. registered under EPA Reg. No. 370 S. Main Street
Goway Company
Yuma, AZ 85364 U.S.A.
(800) 883-1844

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
KEEP OUT OF REACH OF CHILDREN

CAUTION

Causes skin and eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Do not take internally.

STORAGE AND DISPOSAL IN THE U.S.

Do not contaminate water, foodstuffs, feed or seed by storage and disposal.

PESTICIDE STORAGE: Keep bag closed to prevent spills and contamination.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

CONTAINER DISPOSAL: Empty container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned or destroyed. Completely empty container into transfer or application equipment. Do not reuse container. Dispose of empty sack in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(For multiwall bags) Completely empty container into application equipment. Dispose of empty container in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

STORAGE AND DISPOSAL IN CANADA

STORAGE: Keep Avadex MicroActive herbicide bag closed to prevent spills and contamination.

Avoid contamination of seed, feed and foodstuffs.

DISPOSAL: Do not reuse this container for any purpose. Thoroughly empty the contents of the container into the application device. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements.

For information on the disposal of unused, unwanted product, contact the manufacturer or the Provincial Regulatory Agency. Contact the manufacturer and the Provincial Regulatory agency in case of a spill, and for clean-up spills.

In the United States: It is a violation of United States law to use this product without having obtained the United States label at the time of purchase and following the U.S. label at the time of application.

In Canada: NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offense under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

10163-287

It is a violation of United States law to use this product in the United States in a manner inconsistent with its United States labeling.

Far-GO[®] Granular

Herbicide

Granular selective herbicide for fall or spring application to control wild oats in winter wheat, spring and durum wheat, barley, triticale, peas (green, field dried, chickpeas, garbanzo beans), lentils, and sugar beet; and for fall application to suppress *Bromus* species (*B. tectorum*, *B. secalinus* and *B. japonicus*) in winter wheat and in winter barley.

For use in California*, Colorado, Idaho, Kansas, Minnesota, Montana, Nebraska, Nevada, North Dakota, Oregon, South Dakota, Utah, Washington, and Wyoming.

* Not for use on sugar beets in California.

ACTIVE INGREDIENT:

Triallate, S-(2,3,3-trichloroallyl)-diisopropylthiocarbamate

% By Wt.

10.0%

OTHER INGREDIENTS:

90.0%

TOTAL 100.0%

Attention, contains the allergen soy.

KEEP OUT OF REACH OF CHILDREN CAUTION

Read the entire label before using this product. Use only according to label instructions. Read "NOTICE OF CONDITIONS OF SALE AND WARRANTY LIABILITY LIMITATIONS" before buying or using. If terms are not acceptable, return at once unopened.

THIS IS AN END-USE PRODUCT. GOWAN COMPANY DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. SEE INDIVIDUAL CONTAINER LABEL FOR REPACKAGING LIMITATIONS.

FIRST AID Herbicide	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
When seeking medical attention, take the container label if at all possible. If not, take information which identifies the product, that is, the product names and registration numbers.. You may also contact 1-888-478-0798 for emergency medical treatment information.	

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes Moderate Eye Irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

NET CONTENTS _____

EPA Reg. No. 10163-287

EPA Est. No. _____

Gowan
The Go To Company

ACCEPTED
JAN 31 2007

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

10163-287

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Loaders, applicators*, flaggers* and other handlers must wear: Long-sleeved shirt and long pants, shoes plus socks.

A NIOSH approved particulate filtering respirator equipped with N, R, or P class filter media. The respirator should have a NIOSH approval number prefix TC-84A.

* See engineering controls below for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENGINEERING CONTROLS

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for Agricultural Pesticides [40 CFR 170.240(d)(6)].

Flaggers must be in an enclosed cab that meets the definition in the Worker Protection Standard for Agricultural Pesticides for dermal protection and in addition to wearing the required PPE specified above, have immediately available for use in case they must leave the cab: coveralls, chemical-resistant gloves and chemical-resistant footwear, such as nitrile rubber, neoprene, neoprene rubber or polyethylene.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

Ground Water Advisory

Triallate has a degradation product TCPSPA (trichloropropene sulfonic acid) with properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Surface Water Advisory

Under some conditions, the triallate degradate TCPSPA may have a high potential for runoff into surface water (primarily via dissolution in runoff water). These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas over-laying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain surface water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product can only be used in accordance with the Directions for Use on this label or in separately published Gowan Company Supplemental Labeling.

Do not allow this product to drift. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete following application.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: if the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves, such as nitrile rubber, neoprene rubber or polyethylene.

For more options, follow the instructions for category A (dry and water-based formulations) on an EPA chemical-resistant category selection chart.

GENERAL INFORMATION

This herbicide is recommended for wild oat control in barley, peas (green, field dried, chickpeas, garbanzo beans), lentils, durum, spring and winter wheat, triticale, and sugar beet only, and for suppression of downy brome (*Bromus tectorum*), cheat (*Bromus secalinus*) and Japanese brome (*Bromus japonicus*) in winter wheat and winter barley. Other crops should not be treated with this product because injury may occur. For barley, durum, spring wheat, triticale and sugar beet, this product may be applied on the soil either in the fall or in the spring before wild oats germinate. For lentils and peas, this product may only be applied in the spring before wild oats germinate. For winter wheat, this product may only be applied in the fall before wild oats germinate.

Application equipment must be properly calibrated: application of too much herbicide may injure the crop; application of too little may result in poor wild oat control. Recommended rates should be followed in order to:

1. Avoid crop injury
2. Avoid crop residue at harvest
3. Control wild oats
4. Suppress *Bromus* species

GENERAL PRECAUTIONS AND RESTRICTIONS

Application to a field which is wet, lumpy, rough or ridged will result in reduced wild oat control and promote crop thinning. Incorporation must be completed within 48 hours after application and before germination of the wild oats. If weeds, including wild oats, have emerged prior to treating and/or planting, they must be controlled. All deep tillage by cultivators or double disc implements must be completed prior to application. Do not plow following application of this herbicide.

Seeding may be done either before or after application, depending upon the crop that is to be sown. If seeding is delayed, shallow reworking of the treated area before seeding will not destroy the effects of the chemical.

When using this herbicide, a strip should be left untreated for proof of results. Weed control may be evaluated also by removing a surface inch or two inches of the soil at the time of germination to inspect the number of wild oats that were killed before emergence.

Wild oats are usually killed before emergence, but occasionally, and particularly under dry conditions, plants may reach the 3-4 leaf stage before they die.

Under conditions of prolonged high temperature at the time of germination, or extreme drought in the spring, this product may not maintain the usual high standard of wild oat control.

ATTENTION

DELAYED EMERGENCE, STAND REDUCTION, STUNTING AND YIELD LOSS MAY RESULT DUE TO COLD OR WET CONDITIONS, IRRIGATION DURING GERMINATION OR EMERGENCE, DEEP PLANTING, SOIL CRUSTING, DISEASE, INSECTS, OR SELECTION OF A VARIETY GENETICALLY SUSCEPTIBLE TO STRESS. THESE ARE CONDITIONS BEYOND THE CONTROL OF GOWAN COMPANY, AND THE GROWER SHOULD CONSULT THE STATE UNIVERSITY EXTENSION SPECIALIST OR SEED PRODUCER OR SUPPLIER FOR LOCAL VARIETAL INFORMATION AND RECOMMENDED TILLAGE AND PLANTING PRACTICES AND DATES.

LATE PLANTING OF WINTER WHEAT OR PLANTING WINTER WHEAT UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS THAT RESULT IN FAILURE OF THE WHEAT TO EMERGE IN THE FALL MAY RESULT IN STAND REDUCTION, STUNTING AND YIELD LOSS.

Domestic oats should not be seeded where this product was used the previous year.

Do not rotate to crops other than winter wheat, spring and durum wheat, triticale, barley, peas (green, field dried, chickpeas, and garbanzo beans), lentils, and sugar beets for 12 months after a Far-Go application.

Do not graze livestock on treated crops.

When this product is used on clay knobs, crop thinning may occur due to uneven seeding depth and poor crop establishment conditions. Thinning is usually offset by increased tillering and by the reduction of wild oat competition.

IN THE STATE OF MONTANA, do not use this product on fields to be seeded to hard red spring wheat with press drills, if the field is or will be irrigated in the current growing season.

Do not use post-plant incorporation with hoe drills. This practice will result in unacceptable seeding depth and an excessive concentration of this product in the seed zone.

Use of this product not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. Keep container closed to prevent spills and contamination.

APPLICATION EQUIPMENT AND TECHNIQUES

This product must be applied through a specially designed ground applicator or airplane capable of applying small quantities of granules evenly.

Application is limited to one per growing season and must not exceed 15 pounds of Granular Far-Go herbicide per acre.

GROUND EQUIPMENT:

It is important that the applicator be calibrated properly to deliver the desired amount of this product to avoid applying too little, or too much, material. To give even distribution, scatter plates (similar to those used for applying granules in a band) must be attached to each delivery tube or outlet in such a manner to give overall coverage. To calibrate, attach a collector pan, or bag, over each spreader plate or delivery tube. Operate over normal terrain to be treated at 4 to 5 miles per hour. Collect the granules from all outputs after covering the desired distance.

For example: If a 14-foot applicator is being used, stake off a distance of 324 feet in the field to be treated (distance equal to 1/10 acre treated). After attaching a bag or a collection pan to each outlet, collect the granules while driving the desired distance. Check to see that each outlet disperses the same amount of granules. Combine all samples and weigh. For the above set of conditions, the quantity of granules that should be collected for the following recommended rates of broadcast treatment are:

APPLICATION RATES PER ACRE

	<u>10 lbs.</u>	<u>12.5 lbs.</u>	<u>15 lbs.</u>
Amount to be collected:	1 lb.	1.25 lbs.	1.5 lbs.

If more or less than the desired quantity is collected, adjust accordingly and again collect the granules while driving the staked-off distance. Continue this procedure until the required amount is delivered.

AIRPLANE: For aerial application, attachments designed for applying low volumes of granules must be used. In order to ensure uniform aerial application, it is recommended that the field distribution pattern is checked and any necessary gate and vent modifications are made to ensure an even pattern distribution. In order to ensure uniform application and to avoid overlapping and possible crop injury, it is recommended that two flaggers, one at each end of the field, be used.

FIELD PREPARATION

Before applying this product, be sure the soil is in good working condition. All deep tillage by cultivation, or double disc implements, must be completed prior to application of this product. If stubble ground is being treated, one or two passes with a field cultivator may be required before application and incorporation.

INCORPORATED APPLICATIONS

When summer fallow or plowed ground is being treated, a field cultivator will provide adequate incorporation. For proper incorporation of this product, set incorporation implement to work the soil no deeper than 3 to 4 inches. Do not use disc implements for incorporation.

FALL - On summer fallow ground which is loose and free of lumps and trash, incorporate this product using equipment similar to a culti-harrow or a duck-foot with rod weeder attachment.

For applications to fields or standing stubble, fields may be worked once or twice with a field cultivator or chisel plow. Apply granules and incorporate with a field cultivator or culti-harrow.

If soil must be ridged after incorporating to prevent soil erosion by high winds, the depth of ridging should be kept to a minimum.

In preparing the herbicide-treated area for seeding in the spring, care must be taken to avoid working the soil any deeper than fall tillage.

When incorporating with a single pass in the fall, a second incorporation must be performed in the spring during seedbed preparation. If no spring work is anticipated, then both incorporations should be done in the fall.

For applications to winter wheat after seeding, apply and shallowly incorporate with a spike tooth, or spring tooth harrow set shallowly as not to disturb wheat seed.

SPRING – Before Seeding Incorporation: For applications to fields of standing stubble, work fields once or twice with a disc, field cultivator or chisel plow, to provide soil in a good working condition. Apply granules and incorporate with equipment such as a culti-harrow or duck-foot cultivator. A second incorporation at right angles should provide best results. A delay of at least 3 days between the first and second incorporation is recommended for optimum performance.

FOR SPRING AND DURUM WHEAT AND BARLEY IN MONTANA ONLY – For suppression of Persian damel (*Lolium temulentum*), apply this product before seeding and shallowly incorporate.

After Seeding Incorporation: Apply granules immediately after seeding and shallowly incorporate at right angles with equipment such as a flex multi-weeder or harrow. Adjust incorporation equipment to a depth so as not to disturb the seed.

FALL TREATMENT INCORPORATED

CROP	RATE (lb/A)	WHEN TO APPLY
Spring and Durum Wheat, and Barley	12.5 to 15	*Within 3 weeks of normal freeze-up, or until snow cover occurs.
Winter Wheat and Winter Barley**†	12.5 to 15	Before or after seeding.
Sugar beet (Except CA)	15	Fall application – within three weeks of normal freeze-up or until snow cover occurs.

Apply lower rates on light soils and apply higher rates on heavy soils.

Incorporate within 48 hours. For methods, see "FIELD PREPARATION" and "INCORPORATED APPLICATIONS" sections of this label.

* Breakdown of this product is minimal at soil temperatures of 40°F or less.

** For suppression of downy brome (*Bromus tectorum*), cheat (*Bromus secalinus*) and Japanese brome (*Bromus japonicus*) apply 15 pounds per acre prior to planting and shallowly incorporate. In Colorado, Kansas, Nebraska and South Dakota plant with hoeddrills only.

† For winter wheat in the Pacific Northwest, some crop thinning may occur, especially on clay knobs, where seed is "dusted in" or "floated on" and emergence is delayed. Thinning is usually more than offset by tillering and increased yields.

SPRING TREATMENT INCORPORATED

CROP	RATE (lbs/A)	WHEN TO APPLY
Spring and Durum Wheat	10 to 12.5	After seeding before wild oats germinate.
Spring and Durum Wheat	10	Before seeding and before wild oats germinate - Disc/Press drills.
Spring and Durum Wheat	10 to 12.5	Before seeding and before wild oats germinate – Hoeddrills.
Barley and Triticale	12.5 to 15	Just before or immediately after seeding, before wild oats germinate.
Lentils	12.5 to 15	Just before or immediately after seeding, before wild oats germinate.
Peas (green, field dried, chickpeas, garbanzo beans)	12.5 to 15	Just before or immediately after seeding, before wild oats germinate.
Sugar beet (Except CA)	15	Before seeding and before wild oats germinate.

*Apply lower rates on light soils and apply higher rates on heavy soils.

Incorporate within 48 hours. For methods, see "FIELD PREPARATION" and "INCORPORATED APPLICATIONS" sections of this label.

Best spring and durum wheat results will be obtained with at least a 3-day delay between herbicide incorporation and seeding.

SURFACE APPLICATION WITH DELAYED INCORPORATION USE ONLY IN CALIFORNIA, IDAHO, MINNESOTA, MONTANA, NORTH DAKOTA AND UTAH

This product may be applied as a surface application with a delayed incorporation prior to planting spring wheat, durum wheat, and barley at a rate of 15 pounds per acre for all soil types.

Surface applications of this product which will be followed by a delayed incorporation in the spring may be made beginning 3 weeks prior to soil freeze-up in the fall and ending before spring thaw. (Average soil temperature at the 2-inch depth must be 40°F or less).

Surface applications must be incorporated in the spring; two passes are recommended. If surface applications are made because soil conditions do not permit incorporation and these conditions change, making incorporation possible, incorporate even if several weeks after application.

This surface application may be made in standing stubble or to fields with surface residues. Surface applications should not be made to fields covered with snow or with excessive crop residue, which will not allow granule contact with the soil.

Surface applications with delayed incorporation may not provide the same level of wild oat control as fall incorporated applications. Wild oat control resulting from delayed incorporation may be reduced if drought or abnormally warm temperatures occur between application and crop

emergence in the spring. To achieve the most reliable wild oat control, this product should be incorporated within 48 hours of application as described in the "FIELD PREPARATIONS" and "INCORPORATED APPLICATIONS" sections of this label.

NO-TILL APPLICATION FOR WILD OAT CONTROL IN SPRING PLANTED CROPS

This product may be applied prior to planting no-till spring wheat, durum wheat, and barley at a rate of 15 pounds per acre for all soil types.

For best results no-till applications of this product may be made 3 weeks prior to soil freeze-up and ending before spring thaw (average soil temperature at the 2-inch depth must be 40°F or less). Spring applications can also be made if applications are made at least 10 to 14 days prior to seeding.

No-till applications may be made in standing stubble or to fields with surface residue. No-till applications should not be made under conditions which will not allow granule contact with the soil.

No-till applications are to be made in reduced tillage/no-till systems only. Best results should be expected when utilizing drills equipped with high disturbance openers. When seeding with drills equipped with low disturbance openers, or with spring applications, use of heavy harrows or similar minimal disturbance implements prior to seeding may improve control. It is recommended that seeding be delayed until fields begin to green up in the spring and an application of a glyphosate herbicide has been made to control emerged weeds. If this product is used in any other system, it must be applied and incorporated as described elsewhere in this label.

SURFACE APPLICATION FOR FALL PLANTED CROPS

FOR USE IN CALIFORNIA, COLORADO, KANSAS, NEBRASKA AND SOUTH DAKOTA-- This product can be surface-applied prior to seeding of winter wheat without incorporation when seeding is done with a hoeddrill. Apply 15 pounds per acre for suppression of downy brome (*Bromus tectorum*), cheat (*Bromus secalinus*) and Japanese brome (*Bromus japonicus*).

FOR USE IN CALIFORNIA, IDAHO, MONTANA, OREGON, UTAH and WASHINGTON -- Apply this product at a rate of 15 pounds per acre as a surface application ahead of a Yielder™ type no-till drill to obtain suppression of wild oats and downy brome in winter wheat or barley. A flex-tine harrow attached to the drill or used shallowly following planting at a right angle to the drill rows may improve herbicide performance.

If you have any questions concerning this label pamphlet or the proper application or use of Far-GO® Granular Herbicide, contact your local Gowan Company Sales Representative.

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage and disposal.

PESTICIDE STORAGE: Keep bag closed to prevent spills and contamination.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

CONTAINER DISPOSAL: Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned or destroyed. Completely empty container into transfer or application equipment. Do not reuse container. Dispose of empty sack in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

FOR BULK AND MINIBULK CONTAINERS:

Instructions for Users

When the container is empty, replace the cap and seal all openings that have been opened during use; and return the container to the point of purchase, or to an alternate location designated by the registrant at the time of purchase of this product. If not returned to the point of purchase or to a designated location, triple rinse or pressure rinse the empty container and offer for recycling if available.

Instructions for Users and Refillers

This container must only be refilled with this pesticide product.

Do Not Reuse the Container for Any Other Purpose. Do not transport if this container is damaged or leaking. If the container is damaged, leaking, or obsolete, or obtain information about recycling refillable containers, Contact Gowan Company at (800) 883-1844. Cleaning is not necessary prior to refilling with the same product. Clean container before final disposal. Disposal of this container must be in compliance with state and local regulations.

Instructions for Refillers

Prior to refilling inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transporting. If the container cannot be refilled, triple rinse or pressure rinse the empty container and offer for recycling if available.

For Multiwall Bags: Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK OR FIRE), CALL CHEMTREC®(800) 424-9300.

For other product information, contact Gowan Company or see Material Safety Data Sheet.

NOTICE OF CONDITIONS OF SALE AND WARRANTY AND LIABILITY LIMITATIONS

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our recommendations for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of Gowan Company. To the extent consistent with applicable law, all such risks are assumed by the Buyer and User.

Gowan Company warrants that this product conforms to the specifications on the label when used in strict conformance with Direction for Use, subject to the above stated risk limitations. GOWAN COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE FULLEST EXTENT PERMITTED BY LAW, GOWAN COMPANY'S EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT GOWAN COMPANY'S SOLE DISCRETION.

Far-Go® is a registered trademark of Gowan Company LLC.

Roundup® is a registered trademark of Monsanto Technology LLC.

Yielder™ is a trademark of Yielder Drill Company

Far-Go Granular NAFTA label Pending: (to EPA 1-31-07)